

Environmentally important interactions between neptunium(V) and manganese oxide hydroxide minerals

P.A. Wilk¹, D.A. Shaughnessy², C.H. Booth¹, and H. Nitsche^{1,3}

(1) Lawrence Berkeley National Laboratory, Berkeley, CA.

(2) Lawrence Livermore National Laboratory, Livermore, CA.

(3) Department of Chemistry, University of California, Berkeley, CA.

To further understand the fundamental chemical processes governing the migration of transuranic contaminants in the environment, we are currently investigating the interactions between neptunium and manganese oxide/hydroxides. Manganese oxides, present as minor phases in the vadose zone, have been previously shown to sequester large quantities of plutonium under environmental conditions [1,2]. We are now continuing these studies with neptunium(V). Sorption onto manganite and hausmannite has been studied as a function of neptunium concentration and pH. XAFS spectra taken at the neptunium L_{III} edge, have been used to determine the oxidation state of the interacting neptunium as well as a structural probe. Neptunium is expected to sorb to these mineral surfaces in an inner-sphere coordination consistent with other actinides. Our preliminary experimental results indicate interesting redox behavior in this system.

[1] Dawn Shaughnessy, *et al.*, Complexation and Redox Interactions Between Aqueous Plutonium and Manganese Oxide Interfaces, Actinides-2001 proceedings, in press.

[2] D.A. Shaughnessy, *et al.*, Molecular Interfacial Reactions Between Pu(VI) and Manganese Oxide Minerals Manganite and Hausmannite, submitted to Environmental Science and Technology.

Acknowledgement

Supported by the Office of Science and Technology, DOE, Environmental Management Science Program, and by the Office of Basic Energy Sciences, Chemical Sciences, Geosciences, and Biosciences Division. LBNL is operated under the DOE Contract Number DE-AC03-76SF00098.